


POCKET GUIDE: TAPERING OPIOIDS FOR CHRONIC PAIN*

CCN(CC)C(=O)c1ccc(cc1)[C@H](c2ccccc2)N3CC[C@H](C)CC3


Follow up regularly with patients to determine whether opioids are meeting treatment goals and whether opioids can be reduced to lower dosage or discontinued.



GUIDELINE FOR PRESCRIBING OPIOIDS FOR CHRONIC PAIN

*Recommendations focus on pain lasting longer than 3 months or past the time of normal tissue healing, outside of active cancer treatment, palliative care, and end-of-life care.

WHEN TO TAPER



Consider tapering to a reduced opioid dosage or tapering and discontinuing opioid therapy when your patient:

- requests dosage reduction
- does not have clinically meaningful improvement in pain and function (e.g., at least 30% improvement on the 3-item PEG scale)
- is on dosages ≥ 50 MME*/day without benefit or opioids are combined with benzodiazepines
- shows signs of substance use disorder (e.g. work or family problems related to opioid use, difficulty controlling use)
- experiences overdose or other serious adverse event
- shows early warning signs for overdose risk such as confusion, sedation, or slurred speech

*morphine milligram equivalents

HOW TO TAPER

Tapering plans should be individualized and should minimize symptoms of opioid withdrawal while maximizing pain treatment with nonpharmacologic therapies and nonopioid medications. In general:



Go Slow

A decrease of 10% of the original dose per week is a reasonable starting point. Some patients who have taken opioids for a long time might find even slower tapers (e.g., 10% per month) easier.

Discuss the increased risk for overdose if patients quickly return to a previously prescribed higher dose.



Consult

Coordinate with specialists and treatment experts as needed—especially for patients at high risk of harm such as pregnant women or patients with an opioid use disorder.

Use extra caution during pregnancy due to possible risk to the pregnant patient and to the fetus if the patient goes into withdrawal.



Support

Make sure patients receive appropriate psychosocial support. If needed, work with mental health providers, arrange for treatment of opioid use disorder, and offer naloxone for overdose prevention.

Watch for signs of anxiety, depression, and opioid use disorder during the taper and offer support or referral as needed.




Encourage

Let patients know that most people have improved function without worse pain after tapering opioids. Some patients even have improved pain after a taper, even though pain might briefly get worse at first.

Tell patients “I know you can do this” or “I’ll stick by you through this.”

CONSIDERATIONS

**Adjust
Monitor
Reduce**



- 1 Adjust the rate and duration of the taper according to the patient's response.
- 2 Don't reverse the taper; however, the rate may be slowed or paused while monitoring and managing withdrawal symptoms.
- 3 Once the smallest available dose is reached, the interval between doses can be extended and opioids may be stopped when taken less than once a day.

RESOURCES:

CDC Guideline for Prescribing Opioids for Chronic Pain

www.cdc.gov/drugoverdose/prescribing/guideline.html

Washington State Opioid Taper Plan Calculator

www.agencymeddirectors.wa.gov/Files/2015AMDGOpioidGuideline.pdf

Tapering Long-Term Opioid Therapy in Chronic Noncancer Pain

[www.mayoclinicproceedings.org/article/S0025-6196\(15\)00303-1/fulltext](http://www.mayoclinicproceedings.org/article/S0025-6196(15)00303-1/fulltext)



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

www.cdc.gov/drugoverdose